

Product Information Sheet

C1888 Carbendazim

Synonyms: BCM, Methyl 2-benzimidazolecarbamate, Methyl benzimidazol-2-

vlcarbamate

CAS: 10605-21-7 $C_9H_9N_3O_2$ Formula: Mol. Weight: 191.19

Properties

Form: Powder

Appearance: White to Off-White

Solubility: Solubility per the Merck Index in water is 8 mg/mL at pH 7, 29

mg/mL at pH 4. It is also soluble in HCl and Acetic Acid

Plant Tissue Culture Antimycotics Application:

Storage Temp: 2 to 6 °C Stock Solution 2 to 6 °C Storage Temp:

Typical Working

N/A Concentration:

Other Notes: Plant Tissue Culture Tested

Application Notes

Carbendazim is a broad-spectrum systemic antimycotic. Its mode of action is to inhibit the formation of mitotic microtubules in of fungi. In a study, 10 ppm minimum inhibitory concentration of carbendazim is most effective toward of Fusarium mangiferae, a fungus that affects mango plants.² Carbendazim has also been reported to have cytokinin-like activity effect in plant tissue cultures, e.g., Asparagus, Cordyline and Prunus.³

Please Note: It is the sole responsibility of the purchaser to determine the appropriateness of this product for the specific plants that are being cultured and applications that are being used.

References

- 1. Garcia, Pablo C., Rosa M. Rivero, Juan M. Ruiz, and Luid Romero. 2003. The Role of Fungicides in the Physiology of Higher Plants: Implications for Defense Responses. The Botanical Review. 69(2):167-172.
- 2. Iqbal, Zafar, M.A. Pervez, S. Ahmad, Y. Iftikhar, M. Yasin, A. Nawaz, M. Ghazanfar, A. A. Dasti, and A. Saleem. 2010. Determination of minimum inhibitory concentrations of fungicides against fungus Fusariium mangiferae. Pakistan Journal Botany. 42(5). pp. 3525-3532.
- 3. Debergh, P.C., G. De Coster, and W. Steurbaut. 1993. Carbendazim as an Alternative Plant Growth Regulator in Tissue Culture Systems. In Vitro Cellular & Developmental Biology Plant. Vol. 29P. No. 2(April) pp. 89-91.
- 4. Merck Index. 13th Ed. #1799

India Contact