## **PhytoTechnology Laboratories®**

Helping to Build a Better Tomorrow through Plant Science™

### **Product Information Sheet**

M5541

## Murashige & Skoog (MS) Basal Salt Mixture Van der Salm Modification

**Properties** 

Form: Fine Powder

Appearance: Light Burgandy Powder Application: Plant Tissue Culture

Solubility: Water
Typical Working
Concentration: 4.36 g/L

Storage Temp: 2-6°

Storage Temp of Preparation of concentrated solutions is not recommended as insoluble

Stock Solution: precipitates may form.

Other Notes: Contains the macro- and micronutrients as described by Murashige and

Skoog (1962) with the iron source as described by Van der Salm et al.

(1994).

Unadjusted pH = Approx. 3.5-4.5

Formula (mg/L)

Ammonium Nitrate	1650
Boric Acid	6.2
Calcium Chloride, Anhydrous	332.2
Cobalt Chloride•6H <sub>2</sub> O	0.025
Cupric Sulfate•5H <sub>2</sub> O	0.025
FeNa-EDDHA	96.0
Magnesium Sulfate, Anhydrous	180.7

Manganese Sulfate•H <sub>2</sub> O	16.9
Molybdic Acid (Sodium Salt) • 2H <sub>2</sub> O	0.25
Potassium Iodide	0.83
Potassium Nitrate	1900
Potassium Phosphate, Monobasic	170
Zinc Sulfate•7H <sub>2</sub> O	8.6

#### **Application Notes**

Plant Tissue Culture Tested

Plant species: Originally published for use on the rose rootstock cv 'Moneyway' due to chlorosis in shoots cultured on the standard MS formula.

#### References

Murashige, T and F Skoog. 1962. A revised medium for rapid growth and bioassays with tobacco tissue cultures. Physiol. Plant. 15:473-497.

Van der Salm, TMP, CJG Van der Toorn, CH Hänisch ten Cate, LAM Dubois, DP De Vries, and HJM Dons. 1994. Importance of the iron chelate formula for micropropagation of Rosa Hybrida L. 'Moneyway'. Plant Cell Tiss. and Organ Cult. 37:73-77.

Revised 6/2010

### **India Contact**

# Life Technologies (India) Pvt Ltd.